

# AANP Adult Gerontology Primary Care Nurse Practitioner Practice (Sample)

*Study Guide*



*Everything you need from our exam experts!*

**Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.**

**ALL RIGHTS RESERVED.**

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

# Table of Contents

|                             |    |
|-----------------------------|----|
| Copyright .....             | 1  |
| Table of Contents .....     | 2  |
| Introduction .....          | 3  |
| How to Use This Guide ..... | 4  |
| Questions .....             | 5  |
| Answers .....               | 8  |
| Explanations .....          | 10 |
| Next Steps .....            | 16 |

SAMPLE

# Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

# How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

## 1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

## 2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

## 3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

## 4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

## 5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

## 6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

## Questions

SAMPLE

- 1. What common condition necessitates the use of the Beers Criteria in medication management for older adults?**
  - A. Developmental Disabilities
  - B. Chronic Pain
  - C. Polypharmacy
  - D. Depression
- 2. A firm bump measuring 10 mm on a PPD test suggests what?**
  - A. No risk for TB
  - B. TB infection in a low-risk group
  - C. TB infection in a moderate-risk group
  - D. TB infection in a high-risk group
- 3. Which of the following is a characteristic of power of attorney?**
  - A. It allows for the creation of a living will
  - B. It designates someone to make decisions on behalf of another
  - C. It is used for managing healthcare directives only
  - D. It is a document that guarantees medical coverage
- 4. Which organism is NOT one of the three most common bacterial causes of diarrhea?**
  - A. Salmonella
  - B. Campylobacter
  - C. Escherichia coli
  - D. Shigella
- 5. Which medication is indicated for postprandial hyperglycemia?**
  - A. Colesevelam
  - B. Nateglinide
  - C. Glimepiride
  - D. Pioglitazone

- 6. What is a key auscultatory finding of aortic stenosis?**
- A. Rumbling opening snap
  - B. High pitched blowing murmur
  - C. Midsystolic ejection radiating to neck
  - D. Low pitched murmur during diastole
- 7. What should be monitored to detect hypoglycemia in patients taking sulfonylureas?**
- A. Blood glucose levels
  - B. Weight changes
  - C. Liver enzymes
  - D. Cholesterol levels
- 8. How long before a dental procedure should prophylaxis be administered?**
- A. 30 minutes
  - B. 1 hour
  - C. 2 hours
  - D. 24 hours
- 9. What is a classic laboratory finding in hypothyroidism?**
- A. Low TSH and high free T4
  - B. High TSH and low free T4
  - C. Normal TSH and normal free T4
  - D. High TSH and high free T4
- 10. What is the primary purpose of advance care planning?**
- A. To reduce healthcare costs
  - B. To allow individuals to express their wishes regarding future healthcare decisions
  - C. To ensure immediate family consent for treatment
  - D. To prescribe medication ahead of time



## **Answers**

SAMPLE

1. C
2. C
3. B
4. C
5. B
6. C
7. A
8. B
9. B
10. B

SAMPLE

## **Explanations**

SAMPLE

**1. What common condition necessitates the use of the Beers Criteria in medication management for older adults?**

- A. Developmental Disabilities
- B. Chronic Pain
- C. Polypharmacy**
- D. Depression

*The Beers Criteria is a widely recognized guideline that helps healthcare providers avoid prescribing potentially inappropriate medications to older adults. The primary reason for utilizing the Beers Criteria is to address polypharmacy, which refers to the use of multiple medications by a patient, often prevalent in elderly populations due to the multiple chronic conditions they might face. Older adults are particularly vulnerable to the adverse effects of medications, including increased risk of falls, cognitive impairment, and drug interactions. The Beers Criteria serves as a tool for clinicians to identify medications that are high-risk for this demographic, helping to optimize medication regimens and minimize potential harm. This systematic approach supports the safe management of medications, particularly in the context of polypharmacy, where the risk of adverse effects and drug-drug interactions can escalate. By following the Beers Criteria, healthcare providers can provide better care to older adults by ensuring that their medication management aligns with best practices aimed specifically at this population's unique needs.*

**2. A firm bump measuring 10 mm on a PPD test suggests what?**

- A. No risk for TB
- B. TB infection in a low-risk group
- C. TB infection in a moderate-risk group**
- D. TB infection in a high-risk group

*A firm bump measuring 10 mm on a purified protein derivative (PPD) test, commonly used for tuberculosis (TB) screening, indicates a positive reaction, which can suggest an infection with TB. The interpretation of the size of the induration (bump) is influenced by the individual's risk factors for TB. For individuals classified within a moderate-risk group (including those who may have recent exposure to an infectious TB patient or certain medical conditions that slightly increase the risk), an induration of 10 mm or greater is considered a positive test result, signifying the presence of TB infection. This threshold is based on public health guidelines that stratify risk levels and their corresponding induration measurements for interpretation. This finding does not apply to low-risk groups, where the criteria for a positive result is higher (generally requiring a measurement greater than 15 mm for individuals with no known risk factors), and also does not correlate with the high-risk category criteria as those generally require a smaller induration to suggest a positive result. Therefore, in the context of the moderate-risk group, a 10 mm induration is significant and warrants further investigation or treatment based on additional clinical factors.*

### 3. Which of the following is a characteristic of power of attorney?

- A. It allows for the creation of a living will
- B. It designates someone to make decisions on behalf of another**
- C. It is used for managing healthcare directives only
- D. It is a document that guarantees medical coverage

*The characteristic of power of attorney that designates someone to make decisions on behalf of another individual is key to understanding its purpose and function. A power of attorney (POA) allows a person (the principal) to grant legal authority to another individual (the agent or attorney-in-fact) to act on their behalf in specified matters, such as financial, legal, or healthcare decisions. This arrangement is particularly useful in situations where the principal is unable to make decisions for themselves due to illness, incapacity, or absence. The designated agent is empowered to make choices that align with the principal's desires and best interests, effectively ensuring that the principal's preferences are honored even when they cannot articulate them personally. In contrast, other options either misattribute functions to power of attorney or narrow its application too significantly, thus failing to capture the comprehensive nature of what a power of attorney entails. For instance, while a living will is essential in guiding healthcare decisions, it is separate and distinct from a power of attorney. Similarly, power of attorney can encompass various responsibilities beyond healthcare directives, and it does not guarantee medical coverage, as that is the role of insurance or specific healthcare policies.*

### 4. Which organism is NOT one of the three most common bacterial causes of diarrhea?

- A. Salmonella
- B. Campylobacter
- C. Escherichia coli**
- D. Shigella

*Escherichia coli, while a significant pathogen associated with gastrointestinal issues, is often thought of in terms of specific pathogenic strains, such as Enterohemorrhagic E. coli (EHEC), which is linked to outbreaks of hemolytic uremic syndrome (HUS) and severe diarrhea. However, when considering the broader context of bacterial causes of diarrhea, Salmonella, Campylobacter, and Shigella are typically cited as the leading organisms responsible for gastroenteritis in the general population. Salmonella is well-known for causing infections primarily through contaminated food and often leads to severe gastrointestinal symptoms. Campylobacter is frequently associated with undercooked poultry and is the most common bacterial cause of diarrhea in many developed countries. Shigella, on the other hand, is primarily transmitted through contaminated water or food and is notorious for causing dysentery-like symptoms. Thus, while E. coli is indeed a common bacteria, particularly in its pathogenic forms causing illness, it does not rank among the top three overall bacterial causes of diarrhea when considering the more prevalent and widely recognized pathogens. This distinction helps clarify why Escherichia coli is the correct choice in the context of this question.*

## 5. Which medication is indicated for postprandial hyperglycemia?

- A. Colesevelam
- B. Nateglinide**
- C. Glimepiride
- D. Pioglitazone

*Nateglinide is indicated for the management of postprandial hyperglycemia due to its mechanism of action as a rapid-acting insulin secretagogue. This medication stimulates the pancreas to release insulin in response to meals, which is particularly effective in lowering blood sugar levels that rise after eating. This characteristic makes it especially useful in controlling blood glucose spikes that can occur after meals, addressing the postprandial aspect of hyperglycemia. In contrast, the other medications listed have different indications or mechanisms of action that do not specifically target postprandial blood glucose levels. Colesevelam is a bile acid sequestrant used primarily for lowering cholesterol levels and has a secondary role in diabetes management but is not as direct a treatment for postprandial hyperglycemia. Glimepiride, while it also stimulates insulin secretion like nateglinide, has a longer duration of action and is not specifically timed around meals to manage postprandial spikes. Pioglitazone works through improving insulin sensitivity and is primarily effective for fasting hyperglycemia rather than specifically targeting elevations in blood sugar that occur after meals. Thus, nateglinide is the most appropriate choice for addressing postprandial hyperglycemia.*

## 6. What is a key auscultatory finding of aortic stenosis?

- A. Rumbling opening snap
- B. High pitched blowing murmur
- C. Midsystolic ejection radiating to neck**
- D. Low pitched murmur during diastole

*In aortic stenosis, a key auscultatory finding is typically a midsystolic ejection murmur that often radiates to the neck. This murmur occurs due to the turbulent blood flow across the narrowed aortic valve during the systolic phase of the cardiac cycle. The characteristic sound of this murmur is often described as a "crescendo-decrescendo" pattern, which starts softly, peaks, and then diminishes, and it is best heard at the second right intercostal space (the aortic area) and may radiate to the carotid arteries. Because the murmur is associated with the dynamic process of blood ejection from the left ventricle through the stenosed aortic valve, the radiation to the neck is a notable clinical feature. This aspect is crucial for healthcare providers to identify, as it helps in distinguishing aortic stenosis from other types of heart murmurs. In contrast, the other options describe findings associated with different heart conditions. A rumbling opening snap is more indicative of mitral stenosis. A high-pitched blowing murmur is generally associated with aortic regurgitation. A low-pitched murmur during diastole also aligns more with conditions like mitral stenosis or the*

**7. What should be monitored to detect hypoglycemia in patients taking sulfonylureas?**

- A. Blood glucose levels**
- B. Weight changes
- C. Liver enzymes
- D. Cholesterol levels

Monitoring blood glucose levels is essential for detecting hypoglycemia in patients taking sulfonylureas because these medications stimulate the pancreas to release insulin, which lowers blood sugar levels. Patients on sulfonylureas are at risk for hypoglycemia, especially if they do not eat enough carbohydrates, engage in increased physical activity, or take medications that enhance the effects of insulin. Regularly checking blood glucose levels allows for timely identification of low blood sugar and enables prompt intervention to prevent severe hypoglycemic episodes, which can lead to symptoms such as confusion, dizziness, and loss of consciousness. While weight changes, liver enzymes, and cholesterol levels are important indicators in the overall management of diabetic patients, they do not provide direct insight into the immediate risk of hypoglycemia associated with the use of sulfonylureas. Thus, routine monitoring of blood glucose levels stands out as the most crucial practice in this context.

**8. How long before a dental procedure should prophylaxis be administered?**

- A. 30 minutes
- B. 1 hour**
- C. 2 hours
- D. 24 hours

Prophylaxis before a dental procedure is typically recommended to be administered about 1 hour prior to the procedure. This timing allows for the peak therapeutic effect of the antibiotic to occur in the bloodstream, which is crucial for preventing potential bacterial endocarditis in patients at risk. This preventive measure is especially important for individuals with certain heart conditions or other predispositions that could lead to complications from dental procedures that may introduce bacteria into the bloodstream. While administering prophylaxis too soon, such as 30 minutes prior, may not allow enough time for the medication to reach effective blood levels, waiting until 2 hours post-procedure could also increase the risk of infection if bacteria were to enter the bloodstream during the procedure. Lastly, a 24-hour timeframe is excessive for prophylaxis and might not offer any additional benefit. Thus, administering prophylaxis around 1 hour before a dental procedure is considered the optimal approach for effective prevention.

**9. What is a classic laboratory finding in hypothyroidism?**

- A. Low TSH and high free T4
- B. High TSH and low free T4**
- C. Normal TSH and normal free T4
- D. High TSH and high free T4

In the context of hypothyroidism, the classic laboratory finding is characterized by high levels of thyroid-stimulating hormone (TSH) and low levels of free thyroxine (free T4). This occurs because, in hypothyroidism, the thyroid gland is underactive and fails to produce adequate amounts of thyroid hormones. As a result, the pituitary gland responds by increasing the secretion of TSH in an attempt to stimulate the thyroid to produce more hormones. High TSH levels indicate that the body is signaling for more thyroid hormone production due to the low levels of circulating free T4. The low free T4 confirms that thyroid hormone production is insufficient, aligning with the diagnosis of hypothyroidism. This feedback loop between the pituitary gland and the thyroid gland is essential for understanding thyroid function and disorders. In summary, the combination of high TSH and low free T4 is a hallmark of hypothyroidism, validating the choice as the correct answer.

## 10. What is the primary purpose of advance care planning?

- A. To reduce healthcare costs
- B. To allow individuals to express their wishes regarding future healthcare decisions**
- C. To ensure immediate family consent for treatment
- D. To prescribe medication ahead of time

*The primary purpose of advance care planning is to allow individuals to express their wishes regarding future healthcare decisions. This process involves discussing and documenting a person's preferences for medical treatment in situations where they may become unable to communicate their wishes due to illness or incapacity. By engaging in advance care planning, individuals can ensure that their values and desires are respected and upheld, guiding healthcare providers and families in making decisions that align with those preferences. Advance care planning fosters clear communication between patients, their families, and healthcare providers, which is crucial in delivering patient-centered care. It empowers individuals to take an active role in their healthcare, ensuring that their desires—whether they prefer aggressive treatment, palliative care, or other interventions—are known and taken into account when needed. This ultimately leads to more personalized care that aligns with the patient's values, enhancing their quality of life in potentially difficult situations.*

SAMPLE



## Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at [hello@examzify.com](mailto:hello@examzify.com).

Or visit your dedicated course page for more study tools and resources:

<https://aanp.examzify.com>

We wish you the very best on your exam journey. You've got this!

SAMPLE