

Certified Advanced Alcohol and Drug Counselor (CAADC) Practice Exam (Sample)

Study Guide



Everything you need from our exam experts!

Copyright © 2025 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain from reliable sources accurate, complete, and timely information about this product.

SAMPLE

Questions

SAMPLE

- 1. Which of the following is considered a side effect of MAO inhibitors?**
 - A. Weight loss**
 - B. Dry mouth**
 - C. Sensitivity to light**
 - D. Increased exercise tolerance**
- 2. What type of scales includes 3 specific sub-scales in the MMPI-A?**
 - A. Clinical Scales**
 - B. Harris-Lingoes Sub Scales**
 - C. Si Sub Scales**
 - D. Validity Scales**
- 3. A chronically alcoholic veteran exhibits anger towards his employer and has firearms at home. What should you be concerned about?**
 - A. He may be secretly drinking or using drugs**
 - B. He may be expressing Post Acute Withdrawal Syndrome**
 - C. He may be dangerous to others in his workplace**
 - D. He may be experiencing withdrawal symptoms**
- 4. What is a standard criteria for admission to an inpatient program?**
 - A. A court order**
 - B. A need for 24 hour supervision**
 - C. 3 DUI convictions within 5 years**
 - D. Polysubstance dependence**
- 5. Which medication is used to relieve anxiety and may affect chemicals in the brain?**
 - A. Miltown**
 - B. Prozac**
 - C. Zoloft**
 - D. Wellbutrin**

- 6. What is the primary focus of diagnosing Conversion Disorder?**
- A. Presence of multiple psychological disorders**
 - B. Specific symptoms without identifiable physiological cause**
 - C. Specification of psychological stressor related to symptoms**
 - D. All life changes of the individual leading to symptoms**
- 7. What is the primary focus of the MMPI-A assessment?**
- A. Social relationships and skills**
 - B. Patterns in personality and emotional disorders**
 - C. Neuropsychological functioning**
 - D. Cognitive development**
- 8. What distinguishes achievement tests from ability tests?**
- A. Achievement tests are less related to school subjects**
 - B. Achievement tests assess what a client already knows**
 - C. Ability tests are more related to school subjects**
 - D. Achievement tests are more related to school subjects**
- 9. What professional obligation involves monitoring colleagues' practices?**
- A. Peer supervision**
 - B. Self-regulation**
 - C. Accountability**
 - D. Competence maintenance**
- 10. Which of the following is NOT a type of scale found within the MMPI-A?**
- A. Supplementary Scales**
 - B. Behavioral Scales**
 - C. Harris-Lingoes Sub Scales**
 - D. Clinical Scales**

Answers

SAMPLE

1. B
2. C
3. C
4. B
5. A
6. B
7. B
8. D
9. A
10. B

SAMPLE

Explanations

SAMPLE

1. Which of the following is considered a side effect of MAO inhibitors?

A. Weight loss

B. Dry mouth

C. Sensitivity to light

D. Increased exercise tolerance

Dry mouth is indeed considered a side effect of MAO inhibitors. These medications, which are primarily used to treat depression and anxiety, can affect the body's autonomic nervous system, leading to various side effects, one of which is reduced saliva production. This reduction manifests as dryness in the mouth, which can cause discomfort and impact oral health. Weight loss may occasionally be noted in patients on MAO inhibitors due to changes in appetite or gastrointestinal side effects, but it is not as typical or consistent a side effect as dry mouth. Sensitivity to light is generally not associated with MAO inhibitors. Instead, it is more relevant to other medications or conditions. Lastly, increased exercise tolerance does not typically correlate with the effects of MAO inhibitors; in fact, these medications may lead to fatigue or lethargy in some patients, which could decrease rather than improve exercise tolerance.

2. What type of scales includes 3 specific sub-scales in the MMPI-A?

A. Clinical Scales

B. Harris-Lingoes Sub Scales

C. Si Sub Scales

D. Validity Scales

The correct choice identifies the Si Sub Scales as part of the MMPI-A (Minnesota Multiphasic Personality Inventory-Adolescent). These scales focus specifically on areas related to social introversion, helping to assess an adolescent's social functioning and interpersonal relationships. The Si Sub Scales can provide valuable insights into a young person's level of social engagement and are essential for understanding various aspects of their behavioral and emotional functioning. The sub-scales in question are innate to the MMPI-A's framework, which aims to explore personality traits and psychopathology among adolescents. This particular categorization allows clinicians to discern nuances in social behavior and how these might relate to other psychological assessments. Through these insights, practitioners can develop more tailored interventions and support mechanisms for adolescents facing social challenges. Other categories, such as Clinical Scales, Validity Scales, and Harris-Lingoes Sub Scales, serve different functions in assessing the MMPI-A. However, they do not specifically align with the three sub-scales referenced in your query that focus on social introversion.

3. A chronically alcoholic veteran exhibits anger towards his employer and has firearms at home. What should you be concerned about?

- A. He may be secretly drinking or using drugs**
- B. He may be expressing Post Acute Withdrawal Syndrome**
- C. He may be dangerous to others in his workplace**
- D. He may be experiencing withdrawal symptoms**

The situation described involves a chronically alcoholic veteran who shows signs of anger towards his employer and has firearms at home. This scenario raises significant concern for potential violence, particularly in the workplace. Given the factors at play—chronic alcoholism, expressed anger, and access to firearms—there is a heightened risk that this individual could act on their feelings of anger, posing a danger to coworkers or others. Recognizing the risk of violence is crucial, as mental health experts often highlight the correlation between substance abuse, feelings of anger or resentment, and aggressive behavior. In this case, the presence of firearms only amplifies the potential for harmful outcomes, making it essential to consider safety for the workplace environment. While other options may present relevant aspects of the individual's situation, such as potential substance use or withdrawal symptoms, the primary concern should center on the immediate risk to others. Without intervention, the combination of occupational dissatisfaction and accessible weapons can lead to tragic consequences, underscoring the importance of addressing the safety implications promptly.

4. What is a standard criteria for admission to an inpatient program?

- A. A court order**
- B. A need for 24 hour supervision**
- C. 3 DUI convictions within 5 years**
- D. Polysubstance dependence**

The need for 24-hour supervision is a standard criterion for admission to an inpatient program. Inpatient treatment facilities are designed for individuals who require intensive support and monitoring due to the severity of their substance use disorder and associated risks. This level of care is essential for those whose situations could lead to dangerous behaviors, health complications, or withdrawal symptoms that necessitate constant medical oversight and therapeutic intervention. Individuals with significant substance use issues often exhibit conditions that require more than outpatient treatment can provide, and a structured environment that includes around-the-clock supervision can facilitate a safer and more effective recovery process. This criterion ensures that patients have access to immediate assistance and intervention, which can be critical in acute situations. While a court order, multiple DUI convictions, or polysubstance dependence may indicate a need for treatment and could influence the decision for admission, they are not universally recognized criteria for placement in an inpatient setting. The primary focus of inpatient treatment is the level of care needed, particularly the requirement for continuous support and supervision, which makes the need for 24-hour oversight a clear and standard criterion.

5. Which medication is used to relieve anxiety and may affect chemicals in the brain?

- A. Miltown**
- B. Prozac**
- C. Zoloft**
- D. Wellbutrin**

The medication known as Miltown, which is the brand name for meprobamate, is recognized for its use in relieving anxiety. It acts as a tranquilizer and affects various neurotransmitters in the brain, including those associated with anxiety responses. Miltown increases the availability of gamma-aminobutyric acid (GABA), which is an inhibitory neurotransmitter, contributing to its calming effects. In contrast, Prozac and Zoloft are both selective serotonin reuptake inhibitors (SSRIs) used primarily to treat depression but are also effective for anxiety disorders by increasing serotonin levels in the brain. Wellbutrin, on the other hand, is primarily used to treat depression and to help people quit smoking and works on the neurotransmitters dopamine and norepinephrine but is not typically prescribed to relieve anxiety directly. Therefore, Miltown is the choice that specifically aligns with the description of a medication used to relieve anxiety while affecting brain chemicals.

6. What is the primary focus of diagnosing Conversion Disorder?

- A. Presence of multiple psychological disorders**
- B. Specific symptoms without identifiable physiological cause**
- C. Specification of psychological stressor related to symptoms**
- D. All life changes of the individual leading to symptoms**

The primary focus of diagnosing Conversion Disorder is the presence of specific symptoms that cannot be explained by medical or neurological conditions. It is essential to identify these symptoms as manifestations of psychological distress, rather than being directly caused by identifiable physiological issues. While understanding psychological stressors is relevant and can contribute to the overall evaluation and treatment of the disorder, the hallmark of Conversion Disorder lies in the significant physical symptoms that arise in response to the psychological conflict. This distinctive characteristic sets it apart from other psychological and somatic disorders, emphasizing the importance of recognizing the symptoms as a primary aspect during diagnosis. Being able to identify and address these specific symptoms, rather than just looking for the psychological underpinnings or changes in life circumstances, is critical for accurate diagnosis and effective treatment.

7. What is the primary focus of the MMPI-A assessment?

- A. Social relationships and skills
- B. Patterns in personality and emotional disorders**
- C. Neuropsychological functioning
- D. Cognitive development

The primary focus of the MMPI-A (Minnesota Multiphasic Personality Inventory-Adolescent) assessment is on patterns in personality and emotional disorders. This assessment is specifically designed for adolescents and helps to identify a variety of psychological conditions and personality traits. It measures different dimensions of psychological functioning to provide insights into the emotional and behavioral problems that a young person may be experiencing. The MMPI-A consists of several scales that evaluate various psychological conditions, making it a valuable tool for mental health professionals in diagnosing and forming treatment plans for adolescents. By understanding the patterns revealed in the assessment, clinicians can make informed decisions about interventions and therapeutic approaches tailored to the individual's needs. The other options, while relevant to psychological assessments, do not align with the MMPI-A's primary purpose. For example, social relationships and skills may be assessed through other tools, but they are not the main focus of the MMPI-A. Neuropsychological functioning and cognitive development also fall outside the core aim of the MMPI-A, which is centered predominantly on identifying personality and emotional issues.

8. What distinguishes achievement tests from ability tests?

- A. Achievement tests are less related to school subjects
- B. Achievement tests assess what a client already knows
- C. Ability tests are more related to school subjects
- D. Achievement tests are more related to school subjects**

Achievement tests are specifically designed to measure a person's knowledge or skills in specific subjects or area of learning that they have already been exposed to. They evaluate what an individual has learned, often aligning closely with curriculum standards and educational instruction. This makes these tests particularly relevant to school subjects, as they assess a person's proficiency in topics such as mathematics, reading, and science that are typically taught in an educational setting. In contrast, ability tests, also known as aptitude tests, are intended to measure a person's potential to learn or comprehend information in the future. These tests focus on a broader range of cognitive skills, which may not be tied to a specific curriculum or educational experience. Therefore, achievement tests are more directly related to school subjects because they gauge the outcome of past learning experiences, while ability tests assess the potential for future learning regardless of subject matter. This distinction is essential for understanding how different assessments are utilized in educational and counseling settings.

9. What professional obligation involves monitoring colleagues' practices?

- A. Peer supervision**
- B. Self-regulation**
- C. Accountability**
- D. Competence maintenance**

Peer supervision is essential in the field of counseling and addiction treatment as it fosters a collaborative environment among professionals. This obligation involves systematically reviewing and discussing each other's practices, offering constructive feedback, and ensuring adherence to ethical standards. It reinforces best practices and provides support to colleagues, which enhances the overall quality of care provided to clients. In the context of addiction counseling, peer supervision can help identify areas for improvement, recognize potential biases in treatment approaches, and promote continuous professional growth. It creates a platform where counselors can honestly discuss challenges they face with clients and seek advice from their peers, thereby enhancing their clinical skills and effectiveness. This collaborative approach contributes to the professional development of individuals and the integrity of the counseling profession as a whole. By engaging in peer supervision, counselors ensure that they remain accountable to one another and to their clients, ultimately leading to better outcomes in treatment.

10. Which of the following is NOT a type of scale found within the MMPI-A?

- A. Supplementary Scales**
- B. Behavioral Scales**
- C. Harris-Lingoes Sub Scales**
- D. Clinical Scales**

The MMPI-A, or Minnesota Multiphasic Personality Inventory for Adolescents, comprises several distinct scales that are used to assess various psychological constructs. The types of scales included are Clinical Scales, Supplementary Scales, and the Harris-Lingoes Sub Scales, each serving a specific function in evaluating personality and psychopathology. Clinical Scales are designed to assess major psychological conditions and syndromes, providing a foundation for understanding an individual's mental health. Supplementary Scales offer additional insight into other areas not fully captured by the Clinical Scales, enhancing the comprehensiveness of the assessment. The Harris-Lingoes Sub Scales allow for a more nuanced interpretation of specific dimensions within the broader Clinical Scales, helping clinicians identify specific areas of concern. Behavioral Scales, however, are not recognized as one of the categories within the MMPI-A. While behavioral assessments may be relevant in the context of psychological evaluations, they do not form an official part of the MMPI-A's structure. Thus, identifying Behavior Scales as not being part of the MMPI-A aligns with an understanding of the test's design and the specific types of scales it contains. This distinction is crucial for interpreting the assessment results effectively in clinical practice.