


DIPLOMA IN RADIOLOGICAL TECHNOLOGY (DRT) PRACTICE EXAM (SAMPLE) STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

**VISIT US ONLINE FOR
THE FULL VERSION STUDY GUIDE**

<https://radtechdrt.examzify.com>



Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright	1
Table of Contents	2
Introduction	3
How to Use This Guide	4
Questions	5
Answers	8
Explanations	10
Next Steps	16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

Questions

SAMPLE

- 1. What condition presents with abdominal pain and a rigid abdomen?**
 - A. Pyelonephritis
 - B. Ectopic pregnancy
 - C. Placenta previa
 - D. Abruptio placentae
- 2. When comparing glaucoma and cataracts, which statistical test is best to highlight differences?**
 - A. T-test
 - B. Wald test
 - C. Z-test
 - D. F-test
- 3. A patient is prescribed a new medication. Which of the following would require the patient to have regular blood tests to monitor for side effects?**
 - A. ARBs
 - B. ACE inhibitors
 - C. Diuretics
 - D. Statins
- 4. Which stage of lymphoma is localized to a single lymph node or a group in one specific area?**
 - A. Stage III
 - B. Stage IV
 - C. Stage I
 - D. Stage II
- 5. What is the most common gynecologic cancer in the United States?**
 - A. Vaginal
 - B. Uterine
 - C. Cervical
 - D. Ovarian

- 6. What condition is characterized by chest tightness and difficulty completing sentences?**
- A. Pneumonia
 - B. Asthma
 - C. Emphysema
 - D. Acute bronchitis
- 7. Swelling in what area may indicate the presence of a bubo?**
- A. Neck
 - B. Wrist
 - C. Groin
 - D. Knee
- 8. Which bacterium is a common cause of pneumonia in patients with comorbid conditions?**
- A. *Moraxella catarrhalis*
 - B. *Streptococcus pneumoniae*
 - C. *Haemophilus influenzae*
 - D. *Pseudomonas aeruginosa*
- 9. What physical feature may be present in a patient with Klinefelter syndrome?**
- A. Short stature
 - B. Gynaecomastia
 - C. Webbed neck
 - D. Low-set ears
- 10. Which finding is NOT a criterion included in the Dubowitz/Ballard exam to estimate gestational age?**
- A. Thickness and size of breast
 - B. Plantar creases over soles of feet
 - C. Hypertonic flexion of the knees
 - D. Elasticity of cartilage of the nose

Answers

SAMPLE

1. D
2. A
3. B
4. C
5. B
6. B
7. C
8. B
9. B
10. D

SAMPLE

Explanations

SAMPLE

1. What condition presents with abdominal pain and a rigid abdomen?

- A. Pyelonephritis
- B. Ectopic pregnancy
- C. Placenta previa
- D. Abruptio placentae**

The condition characterized by abdominal pain and a rigid abdomen is abruptio placentae. This condition occurs when the placenta detaches from the uterine wall before delivery, leading to significant bleeding and potential complications for both the mother and the fetus. The detachment causes intense abdominal pain and can result in a tight or rigid abdomen due to the accumulation of blood and muscle spasm in response to injury. When evaluating this condition, abdominal rigidity can be associated with the body's response to internal bleeding, as the muscles may tense in reaction to pain or the physical distress caused by the separation of the placenta. In cases of abruptio placentae, this may also include symptoms like vaginal bleeding and fetal distress, depending on the severity of the detachment. The other conditions mentioned generally present with different signs or symptoms that do not align as closely with the combination of abdominal pain and a rigid abdomen. For example, pyelonephritis typically involves flank pain and fever, while ectopic pregnancy may feature unilateral pain and potentially shoulder pain due to diaphragm irritation. Placenta previa usually presents with painless vaginal bleeding rather than abdominal rigidity. Understanding these distinctions helps clarify why abruptio placentae is the most appropriate condition associated with the given symptoms.

2. When comparing glaucoma and cataracts, which statistical test is best to highlight differences?

- A. T-test**
- B. Wald test
- C. Z-test
- D. F-test

The selection of the t-test as the best statistical method to compare glaucoma and cataracts is appropriate when both conditions need to be evaluated concerning their means, especially when dealing with continuous data derived from a smaller sample size. The t-test is designed for comparing the means from two groups to ascertain if they are significantly different from one another. Given that glaucoma and cataracts can exhibit different physical and clinical characteristics, the t-test would effectively assess whether differences in measurements (e.g., intraocular pressure in glaucoma or visual acuity in cataracts) exist between patients with these two conditions. On the other hand, other tests such as the Wald test, Z-test, and F-test serve different purposes. The Wald test is often used in the context of logistic regression or hypothesis testing for coefficients in models, not typically for direct comparisons between two independent groups. The Z-test is similar to the t-test but is generally applicable for larger sample sizes with known population variances, which may not apply in many clinical settings involving specific conditions like glaucoma and cataracts. Lastly, the F-test is predominantly used in the context of variances among groups (like ANOVA) rather than direct mean comparisons between two specific conditions. Thus, the t-test is the most suitable

3. A patient is prescribed a new medication. Which of the following would require the patient to have regular blood tests to monitor for side effects?

A. ARBs

B. ACE inhibitors

C. Diuretics

D. Statins

ACE inhibitors are commonly prescribed medications for managing hypertension and heart failure. These drugs can affect kidney function and electrolyte levels in the body, particularly potassium. Therefore, it is crucial to monitor renal function through regular blood tests, specifically measuring serum creatinine and potassium levels, to detect any potential adverse effects early. While other medications, such as diuretics, also require monitoring for electrolyte imbalances, the frequency and necessity of blood testing is more critical with ACE inhibitors, especially in patients who may be at risk for renal impairment. In contrast, ARBs and statins typically necessitate less frequent blood monitoring under standard therapy conditions unless the patient presents with specific risk factors or complications.

4. Which stage of lymphoma is localized to a single lymph node or a group in one specific area?

A. Stage III

B. Stage IV

C. Stage I

D. Stage II

The correct answer identifies Stage I lymphoma, which signifies that the cancer is confined to a single lymph node or a group of nearby lymph nodes. In this initial stage, the disease has not spread beyond this localized area, making it the most localized form of lymphoma. Understanding lymphoma staging is crucial as it helps determine the extent of the disease and the appropriate treatment options. Stage I indicates earlier and more confined disease, often associated with better prognosis and more treatment options compared to more advanced stages. In contrast, Stages II, III, and IV indicate progressive involvement of additional lymph nodes or organs, with Stage II encompassing two or more lymph nodes on the same side of the diaphragm, Stage III indicating lymphoma that has spread to lymph nodes on both sides of the diaphragm, and Stage IV representing widespread disease affecting organs outside of the lymphatic system. Each subsequent stage implies a greater extent of the disease, which is associated with more complex treatment challenges and variable prognosis.

5. What is the most common gynecologic cancer in the United States?

- A. Vaginal
- B. Uterine**
- C. Cervical
- D. Ovarian

The most common gynecologic cancer in the United States is uterine cancer, specifically endometrial cancer, which arises from the lining of the uterus. Factors contributing to the prevalence of this cancer include hormonal influences, obesity, and age, with the majority of cases diagnosed in women over the age of 50. Uterine cancer is often detected at an early stage due to the symptom of abnormal vaginal bleeding, making it one of the more manageable cancers when caught early. Its incidence has increased in recent years, likely linked to the rise in risk factors such as obesity and an aging population. In contrast, vaginal, cervical, and ovarian cancers, while significant, have lower incidence rates compared to uterine cancer. Cervical cancer rates have declined due to effective screening programs like Pap smears and the HPV vaccination. Ovarian cancer, although highly impactful in terms of mortality, is less common than uterine cancer, presenting more often in the later stages when symptoms are less evident. Vaginal cancer is the rarest among these types, contributing to its lower prevalence rates.

6. What condition is characterized by chest tightness and difficulty completing sentences?

- A. Pneumonia
- B. Asthma**
- C. Emphysema
- D. Acute bronchitis

The condition characterized by chest tightness and difficulty completing sentences is asthma. This respiratory condition affects the airways, causing them to become inflamed and narrowed, leading to difficulty in breathing. When a person with asthma experiences an attack, they may feel tightness in their chest, and the airflow may be restricted, making it hard to take deep breaths. This difficulty directly impacts their ability to speak in complete sentences as the person struggles to get enough air during an episode. Asthma may also present with wheezing, coughing, and shortness of breath, especially during physical activity or exposure to allergens. Understanding the symptoms of asthma is crucial for prompt management and treatment, particularly as they can vary in intensity from mild to severe. Recognizing this condition is essential for the proper care of affected patients in a clinical setting.

7. Swelling in what area may indicate the presence of a bubo?

- A. Neck
- B. Wrist
- C. Groin**
- D. Knee

The presence of a bubo is characterized by the swelling of lymph nodes, particularly in a specific area of the body. When it comes to buboes, these are commonly associated with infections such as bubonic plague or sexually transmitted infections, and they typically occur in the groin, although they can also occur in the axillary (armpit) or cervical (neck) regions depending on the site of the infection or lymphatic drainage. In the case of the groin, genital region, and lower abdomen, the lymphatic vessels drain these areas into the inguinal (groin) lymph nodes. When these lymph nodes become infected or inflamed, they can swell significantly, resulting in the formation of a bubo. This localized swelling is a clinical sign indicating that an infection is present, and it becomes palpably noticeable in the groin area. Swelling in the neck, wrist, or knee might occur for other reasons, such as localized infections, injuries, or inflammatory conditions, but they do not typically indicate the presence of a bubo associated with lymphatic involvement of the groin region. Hence, when considering the characteristic location of a bubo, the groin is the most relevant area for this type of swelling.

8. Which bacterium is a common cause of pneumonia in patients with comorbid conditions?

- A. *Moraxella catarrhalis*
- B. *Streptococcus pneumoniae***
- C. *Haemophilus influenzae*
- D. *Pseudomonas aeruginosa*

Streptococcus pneumoniae is recognized as a leading cause of pneumonia, particularly in patients who have comorbid conditions such as chronic obstructive pulmonary disease (COPD), diabetes, heart disease, or other underlying illnesses. This bacterium is known for its virulence and ability to colonize the respiratory tract, which increases the risk of respiratory infections when the host's immune system is compromised. In individuals with comorbidities, the altered immune response can make them more susceptible to infections, and *Streptococcus pneumoniae* is adept at evading the host's defenses. The presence of these underlying health issues often results in more severe and complicated pneumonia outbreaks, leading to higher rates of hospitalization and treatment challenges. While other bacteria listed can also cause pneumonia, they do not match the frequency with which *Streptococcus pneumoniae* is associated with pneumonia in patients with compromised health. *Moraxella catarrhalis* and *Haemophilus influenzae* typically cause respiratory tract infections but are less common in pneumonias among this specific group. *Pseudomonas aeruginosa* is more commonly associated with pneumonia in patients who are immunocompromised or hospitalized, rather than in those with generalized comorbidities.

9. What physical feature may be present in a patient with Klinefelter syndrome?

- A. Short stature
- B. Gynaecomastia**
- C. Webbed neck
- D. Low-set ears

Klinefelter syndrome is a genetic condition that occurs in males who have an extra X chromosome, typically manifesting as XXY rather than the usual XY. One of the most notable physical features associated with Klinefelter syndrome is gynaecomastia, which refers to the development of breast tissue in males. This occurs due to hormonal imbalances, particularly with increased estrogen levels relative to testosterone, and can be an identifying characteristic of the syndrome. Gynaecomastia can vary in severity and may cause psychological distress or embarrassment for those affected. It is important for healthcare providers to recognize this condition, as it can have implications for managing not just the physical aspect but also the social and psychological impacts on the patient. The other features mentioned, such as short stature, webbed neck, and low-set ears, are not specifically characteristic of Klinefelter syndrome. Short stature can occur in various genetic or health conditions but is not definitive for Klinefelter. Webbed neck is more commonly associated with Turner syndrome, which affects females. Low-set ears can be a feature of multiple syndromes and are not unique to Klinefelter syndrome. Recognizing the hallmark features of Klinefelter syndrome, such as gynaecomastia

10. Which finding is NOT a criterion included in the Dubowitz/Ballard exam to estimate gestational age?

- A. Thickness and size of breast
- B. Plantar creases over soles of feet
- C. Hypertonic flexion of the knees
- D. Elasticity of cartilage of the nose**

The correct answer is related to the criteria used in the Dubowitz/Ballard examination, which assesses gestational age through various physical and neurological characteristics of newborns. The Dubowitz/Ballard exam includes specific features such as the thickness and size of breast tissue, the presence and distribution of plantar creases, and the degree of flexion in the limbs as indicators of gestational age. The elasticity of cartilage in the nose is not a recognized criterion in this specific assessment. While some examinations may assess nose shape or other characteristics, the Dubowitz/Ballard examination primarily focuses on systematically quantifying physical features and neuromuscular maturity that correlate with gestational age. Therefore, while evaluating newborn gestational age, the aspects determined by this exam are concentrated on more direct physical indicators, leaving out characteristics like nasal cartilage elasticity.

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

<https://radtechdrt.examzify.com>

We wish you the very best on your exam journey. You've got this!

SAMPLE