

NAPLEX (NORTH AMERICAN PHARMACIST LICENSURE EXAMINATION) PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR
THE FULL VERSION STUDY GUIDE

<https://naplex.examzify.com>



Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright	1
Table of Contents	2
Introduction	3
How to Use This Guide	4
Questions	5
Answers	8
Explanations	10
Next Steps	15

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

Questions

SAMPLE

- 1. Which opioid allergy commonality is given in the text?**
 - A. Drugs ending in -fen
 - B. Drugs with -COD or -MORPH in name
 - C. Drugs starting with Tram-
 - D. Medications in liquid form
- 2. Which publication is a reference for adverse drug reactions?**
 - A. FDABLE
 - B. ASHP's Drug-Induced Disease: Prevention, Detection, and Management
 - C. Meyler's side effects of drugs
 - D. Safety reporting portal
- 3. What is the main action of PAMORAS on opioid-induced constipation?**
 - A. Increase bulk formation
 - B. Stimulate peristalsis
 - C. Block opioid receptors in the gut
 - D. Reduce intestinal fluid absorption
- 4. What is the frequency for maintaining a pharmacy temperature log for vaccine freezers?**
 - A. Once daily
 - B. Twice daily
 - C. Weekly
 - D. Monthly
- 5. What is a component of progestin-only pills (Mini-pill/POP)?**
 - A. Errin contains ethinylestradiol
 - B. Norethindrone is found in 'Nor' variants like Nora-BE
 - C. Slynd contains a combination of drospirenone and norethindrone
 - D. All mini-pills contain estrogen
- 6. What is a major risk associated with Thiazolidinediones?**
 - A. Hypoglycemia
 - B. Exacerbation of HF
 - C. Vitamin B12 deficiency
 - D. Renal toxicity

7. Omalizumab (Xolair) is indicated for?

- A. Moderate-severe persistent allergic asthma
- B. Chronic obstructive pulmonary disease (COPD)
- C. Mild intermittent asthma
- D. Acute asthma attacks

8. Which antibiotic should be avoided in pregnancy for malaria prophylaxis?

- A. Doxycycline (Doryx, vibramycin)
- B. Chloroquine
- C. Mefloquine
- D. Atovaquone/proguanil (malarone)

9. For uncomplicated UTI in females, how is Fosfomycin administered?

- A. Intravenously
- B. In a single dose
- C. Twice daily for five days
- D. Once daily for seven days

10. What common application do both Venlafaxine and Duloxetine share?

- A. Depression
- B. Insomnia
- C. Epilepsy
- D. High blood pressure

Answers

SAMPLE

1. B
2. B
3. C
4. B
5. B
6. B
7. A
8. A
9. B
10. A

SAMPLE

Explanations

SAMPLE

1. Which opioid allergy commonality is given in the text?

- A. Drugs ending in -fen
- B. Drugs with -COD or -MORPH in name**
- C. Drugs starting with Tram-
- D. Medications in liquid form

The correct answer is B. Drugs with -COD or -MORPH in the name are commonly associated with opioid allergies. Opioid medications like codeine and morphine fall into this category. This commonality is important for healthcare providers to be aware of when managing patients with known or suspected opioid allergies. Options A, C, and D are not accurate commonalities for opioid allergies. Drugs ending in -fen (Option A) may refer to medications like fentanyl, which is an opioid but not necessarily a commonality for opioid allergies. Drugs starting with Tram- (Option C) may be referring to tramadol, another opioid, but not all drugs starting with Tram- are commonly associated with opioid allergies. Medications in liquid form (Option D) are not specifically linked to opioid allergies; the form of the medication does not determine the likelihood of an allergic reaction.

2. Which publication is a reference for adverse drug reactions?

- A. FDable
- B. ASHP's Drug-Induced Disease: Prevention, Detection, and Management**
- C. Meyler's side effects of drugs
- D. Safety reporting portal

ASHP's "Drug-Induced Disease: Prevention, Detection, and Management" is a reference for adverse drug reactions. This publication provides comprehensive information on drug-induced diseases, including how to prevent, detect, and manage them. It is a reliable source for healthcare professionals to understand the relationship between medications and adverse reactions. Option A, "FDable," is not a specific reference publication for adverse drug reactions. Option C, "Meyler's side effects of drugs," is a well-known reference for adverse drug reactions; however, the ASHP publication mentioned in the correct answer is more focused on the prevention, detection, and management of drug-induced diseases. Option D, "Safety reporting portal," is a reporting platform rather than a reference publication for adverse drug reactions.

3. What is the main action of PAMORAS on opioid-induced constipation?

- A. Increase bulk formation
- B. Stimulate peristalsis
- C. Block opioid receptors in the gut**
- D. Reduce intestinal fluid absorption

PAMORAS is a medication used to treat opioid-induced constipation. This means that the main action of PAMORAS is to address constipation caused by the use of opioids. Option A and B are incorrect because they both refer to actions that aim to alleviate constipation in general rather than specifically addressing opioid-induced constipation. Option D is incorrect because it suggests reducing intestinal fluid absorption would address opioid-induced constipation, which is not the case. The correct answer, Option C, refers to blocking opioid receptors in the gut, which is the main mechanism of action of PAMORAS in addressing constipation caused by opioids. By doing so, PAMORAS can help alleviate constipation and improve bowel movements for individuals who use opioids.

4. What is the frequency for maintaining a pharmacy temperature log for vaccine freezers?

- A. Once daily
- B. Twice daily**
- C. Weekly
- D. Monthly

A Once daily - This option is incorrect because maintaining a pharmacy temperature log for vaccine freezers requires more frequent monitoring than just once a day. Vaccines are sensitive to temperature changes and need to be monitored closely to ensure they remain effective. C: Weekly - This option is incorrect because a weekly frequency for maintaining a temperature log is too infrequent. Temperature fluctuations can occur within a week and may not be caught in time to prevent damage to the vaccines. D: Monthly - This option is incorrect because a monthly frequency for temperature log maintenance is too infrequent. Monthly monitoring may not catch temperature fluctuations or issues that may arise between monthly checks. Bimonthly log maintenance is also not ideal as it is still not frequent enough to catch potential problems before they arise. Thus, the correct answer is B: Twice daily. This frequency allows for more frequent monitoring to catch any temperature fluctuations or issues that may arise and maintain the effectiveness of the vaccines.

5. What is a component of progestin-only pills (Mini-pill/POP)?

- A. Errin contains ethinylestradiol
- B. Norethindrone is found in 'Nor' variants like Nora-BE**
- C. Slynd contains a combination of drospirenone and norethindrone
- D. All mini-pills contain estrogen

Progestin-only pills, or mini-pills, do not contain estrogen. Option A, Errin, contains ethinylestradiol which is a form of estrogen. Option C, Slynd, contains a combination of progestin drospirenone and norethindrone, and thus not a progestin-only pill. Option D is incorrect because as stated before, mini-pills do not contain estrogen. Option B, Nora-BE, contains norethindrone, a type of progestin, making it the correct answer.

6. What is a major risk associated with Thiazolidinediones?

- A. Hypoglycemia
- B. Exacerbation of HF**
- C. Vitamin B12 deficiency
- D. Renal toxicity

Thiazolidinediones is a class of medication used to treat type 2 diabetes. One of the major risks associated with it is exacerbation of heart failure (HF). This is because these medications can cause fluid buildup in the body, which can worsen heart failure symptoms. While hypoglycemia (low blood sugar), vitamin B12 deficiency, and renal toxicity are all potential side effects of thiazolidinediones, they are not the major risk associated with them. Hypoglycemia can be managed by carefully monitoring blood sugar levels and adjusting the dosage of the medication, while vitamin B12 deficiency and renal toxicity can be monitored through regular blood tests.

7. Omalizumab (Xolair) is indicated for?

A. Moderate-severe persistent allergic asthma

B. Chronic obstructive pulmonary disease (COPD)

C. Mild intermittent asthma

D. Acute asthma attacks

Omalizumab (Xolair) is indicated for moderate-severe persistent allergic asthma. It is a monoclonal antibody that is specifically approved for use in patients with allergic asthma that is not controlled with inhaled corticosteroids. The drug works by blocking immunoglobulin E (IgE), which is responsible for triggering allergic reactions in the body. Therefore, option A is the correct choice as it accurately reflects the approved indication for omalizumab. Options B, C, and D are incorrect: B. Chronic obstructive pulmonary disease (COPD): Omalizumab is not indicated for COPD as it is specifically approved for allergic asthma. C. Mild intermittent asthma: Omalizumab is used for moderate-severe persistent allergic asthma, not for mild intermittent asthma. D. Acute asthma attacks: Omalizumab is not used for acute asthma attacks, rather it is a long-term controller medication for persistent allergic asthma.

8. Which antibiotic should be avoided in pregnancy for malaria prophylaxis?

A. Doxycycline (Doryx, vibramycin)

B. Chloroquine

C. Mefloquine

D. Atovaquone/proguanil (malarone)

Doxycycline is the only antibiotic in the list of options, therefore, it should be avoided in pregnancy for malaria prophylaxis. The other options are not antibiotics and are used specifically for malaria prevention, not treatment. Chloroquine and Atovaquone/proguanil (malarone) are typically recommended for use during pregnancy, while Mefloquine should be used with caution and only if no other options are available. It is important to consult with a healthcare provider for the best option for individual circumstances.

9. For uncomplicated UTI in females, how is Fosfomycin administered?

A. Intravenously

B. In a single dose

C. Twice daily for five days

D. Once daily for seven days

For uncomplicated UTI in females, Fosfomycin is typically administered in a single dose. This is because Fosfomycin has a unique characteristic of being effective as a single-dose treatment due to its pharmacokinetics and pharmacodynamics. The drug achieves high concentrations in the urine shortly after administration, making it suitable for a one-time dosing regimen to effectively treat uncomplicated UTIs. The other options are incorrect because Fosfomycin is not administered intravenously, and it is not typically given as a twice-daily regimen for five days or a once-daily regimen for seven days for uncomplicated UTIs.

10. What common application do both Venlafaxine and Duloxetine share?

A. Depression

B. Insomnia

C. Epilepsy

D. High blood pressure

Both Venlafaxine and Duloxetine are commonly used as antidepressants in the treatment of depression. They are both classified as serotonin-norepinephrine reuptake inhibitors (SNRIs), which work by increasing the levels of serotonin and norepinephrine in the brain, thus improving mood and reducing symptoms of depression. These medications are not typically used for insomnia, epilepsy, or high blood pressure.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

<https://naplex.examzify.com>

We wish you the very best on your exam journey. You've got this!

SAMPLE