

STROKE CERTIFIED REGISTERED NURSE (SCRN) PRACTICE EXAM (SAMPLE)

STUDY GUIDE



**SAMPLE STUDY GUIDE
WITH LIMITED QUESTIONS**

**VISIT US ONLINE FOR
THE FULL VERSION STUDY GUIDE**

<https://strokecertifiedregisterednurse.examzify.com>



Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright	1
Table of Contents	2
Introduction	3
How to Use This Guide	4
Questions	5
Answers	8
Explanations	10
Next Steps	16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

Questions

SAMPLE

- 1. What adaptive equipment might be recommended for a stroke survivor?**
 - A. Indoors exercise equipment
 - B. Grab bars, reachers, or mobility aids
 - C. Luxury wheelchairs
 - D. Advanced gaming consoles for rehabilitation
- 2. When should rehabilitation services begin for stroke patients?**
 - A. Immediately after the stroke
 - B. As soon as possible, preferably in the acute phase
 - C. Only after hospital discharge
 - D. Within one month post-stroke
- 3. What is a common nutritional recommendation for stroke survivors?**
 - A. Increased intake of sugars
 - B. Adopting a low-sodium diet
 - C. High protein diets only
 - D. Fasting for health benefits
- 4. Which vital sign is crucial to monitor closely after thrombolytic therapy?**
 - A. Temperature
 - B. Blood pressure
 - C. Respiratory rate
 - D. Heart rate
- 5. What syndrome results from damage to the spinothalamic tract in PICA strokes?**
 - A. Locked-in syndrome
 - B. Wallenberg syndrome
 - C. Weber's syndrome
 - D. Horner's syndrome
- 6. Which cognitive change can result from a stroke affecting the frontal lobe?**
 - A. Improved memory retention
 - B. Impaired judgment or decision-making
 - C. Enhanced analytical skills
 - D. Greater attention span

7. What is a common symptom of stroke?

- A. Severe headache
- B. Sudden numbness or weakness of the face, arm, or leg
- C. Chest pain
- D. Difficulty breathing

8. Which change in speech may indicate a patient has had a stroke?

- A. Clear speech with normal volume
- B. Slurred speech or difficulty finding words
- C. Excessive talking
- D. Sudden increase in vocabulary

9. Which condition is primarily indicated by Wallenberg syndrome?

- A. Basilar artery occlusion
- B. PICA stroke
- C. MCA stroke
- D. ACA stroke

10. How can caregivers best support stroke recovery?

- A. By minimizing patient interactions
- B. By providing encouragement and assistance with therapy exercises
- C. By discouraging any physical activity
- D. By focusing solely on medication management

Answers

SAMPLE

1. B
2. B
3. B
4. B
5. B
6. B
7. B
8. B
9. B
10. B

SAMPLE

Explanations

SAMPLE

1. What adaptive equipment might be recommended for a stroke survivor?

- A. Indoors exercise equipment
- B. Grab bars, reachers, or mobility aids**
- C. Luxury wheelchairs
- D. Advanced gaming consoles for rehabilitation

The recommendation of grab bars, reachers, or mobility aids for stroke survivors focuses on enhancing independence and safety in daily activities. After a stroke, individuals often experience varying levels of mobility impairment and may face challenges with strength, coordination, and balance. Adaptive equipment such as grab bars provides stability in areas of the home where falls are more likely, such as bathrooms and stairways. Reachers help individuals pick up items without straining themselves or losing balance, while mobility aids like canes, walkers, or wheelchairs facilitate better movement and accessibility. These tools not only promote safety but also encourage participation in daily activities, helping survivors to regain confidence and improve their quality of life. Other options, while they may be beneficial in specific contexts, do not fit as universally appropriate adaptive equipment for stroke survivors. For instance, indoor exercise equipment may be too advanced for some individuals at first, luxury wheelchairs do not address the functional needs as directly, and advanced gaming consoles might not be the most practical or effective means of rehabilitation compared to more fundamental aids.

2. When should rehabilitation services begin for stroke patients?

- A. Immediately after the stroke
- B. As soon as possible, preferably in the acute phase**
- C. Only after hospital discharge
- D. Within one month post-stroke

Starting rehabilitation services as soon as possible, preferably in the acute phase, is crucial for stroke patients because early intervention can significantly influence outcomes and recovery. Initiating rehabilitation during the acute phase takes advantage of the neuroplasticity of the brain, which is the ability of the brain to reorganize itself by forming new neural connections. This is particularly vital after a stroke when the brain may begin to adapt to new pathways for motor and cognitive functions. Research indicates that beginning rehabilitation early can help improve mobility, reduce disability, and enhance the overall quality of life. Interventions in this time frame may include physical therapy, occupational therapy, and speech-language therapy, which can aid in regaining lost skills and promoting independence. The acute phase is typically the period right after the stroke event when the patient is still hospitalized, making it the optimum time for various rehabilitation strategies to be implemented. Other options suggest starting rehabilitation at later stages, such as only after hospital discharge or within a month post-stroke, which may delay potential recovery benefits. A brief pause in starting rehabilitation can hinder recovery and decrease the effectiveness of treatment strategies that leverage the body's healing processes shortly after a stroke. Therefore, beginning rehabilitation as soon as possible aligns with best practices in stroke recovery and optimal patient outcomes.

3. What is a common nutritional recommendation for stroke survivors?

- A. Increased intake of sugars
- B. Adopting a low-sodium diet**
- C. High protein diets only
- D. Fasting for health benefits

One common nutritional recommendation for stroke survivors is to adopt a low-sodium diet. High sodium intake is associated with increased blood pressure, which is a significant risk factor for stroke. By reducing salt in the diet, stroke survivors can better manage their blood pressure and reduce the risk of subsequent strokes. This recommendation aligns with general cardiovascular health guidelines, which emphasize the importance of a heart-healthy diet in promoting recovery and preventing further cerebrovascular incidents. Other options focus on dietary approaches that are either not appropriate or lack supportive evidence for stroke recovery. For instance, increased intake of sugars can lead to weight gain and worsen metabolic health, while high protein diets without proper balance of other macronutrients may not adequately address overall dietary needs. Fasting for health benefits is not a universally recommended strategy for stroke survivors, as it might lead to nutritional deficiencies or inadequate caloric intake necessary for recovery. Therefore, a low-sodium diet stands out as a well-supported recommendation for enhancing post-stroke health and prevention.

4. Which vital sign is crucial to monitor closely after thrombolytic therapy?

- A. Temperature
- B. Blood pressure**
- C. Respiratory rate
- D. Heart rate

Monitoring blood pressure closely after thrombolytic therapy is crucial because it helps assess the risk of complications, particularly hemorrhage or bleeding, which can significantly occur following such treatment. Thrombolytic therapy works by dissolving blood clots, and while it can restore blood flow, it can also disrupt blood vessel integrity, leading to increased risk of intracranial hemorrhage or other bleeding complications if blood pressure is not appropriately controlled. High blood pressure following thrombolysis can exacerbate this risk, as elevated pressures may stress weakened blood vessels and increase the likelihood of rupture. Therefore, maintaining optimal blood pressure within a target range is essential to minimize complications and ensure the safety of the patient during the recovery phase post-thrombolysis.

5. What syndrome results from damage to the spinothalamic tract in PICA strokes?

- A. Locked-in syndrome
- B. Wallenberg syndrome**
- C. Weber's syndrome
- D. Horner's syndrome

Wallenberg syndrome, also known as lateral medullary syndrome, occurs when there is damage to the spinothalamic tract and other structures in the lateral medulla of the brainstem, particularly due to a posterior inferior cerebellar artery (PICA) stroke. The spinothalamic tract is responsible for carrying pain and temperature sensations from the body to the brain. Damage to this tract can lead to loss of pain and temperature sensation on the contralateral side of the body, while the ipsilateral side may experience other symptoms due to lesions affecting cranial nerves and other nearby tracts. In addition to the sensory deficits, Wallenberg syndrome presents with a variety of additional neurological deficits, such as ataxia, dizziness, dysphagia, and Horner's syndrome due to involvement of the descending sympathetic fibers. Understanding that Wallenberg syndrome results specifically from PICA strokes and the related damage to the spinothalamic tract helps clarify the pathophysiology involved and the clinical picture that arises from this condition. The other syndromes listed are associated with different vascular territories or anatomical lesions, making them separate from the characteristics of Wallenberg syndrome.

6. Which cognitive change can result from a stroke affecting the frontal lobe?

- A. Improved memory retention
- B. Impaired judgment or decision-making**
- C. Enhanced analytical skills
- D. Greater attention span

A stroke affecting the frontal lobe can lead to significant changes in cognition, particularly in judgment and decision-making abilities. The frontal lobe is crucial for executive functions, which include reasoning, planning, problem-solving, and the ability to control impulses. When a stroke impairs this area, individuals may struggle with assessing situations effectively and making sound decisions. This can manifest as poor judgment in daily activities, difficulty weighing options, or an inability to foresee the consequences of actions, ultimately impacting their ability to interact successfully with their environment. In contrast, options like improved memory retention, enhanced analytical skills, and a greater attention span do not typically align with the cognitive deficits observed following frontal lobe damage. These functions may actually decline as executive functions become disrupted in the wake of a stroke. Understanding this relationship between stroke location and cognitive function is essential for providing appropriate care and support to affected individuals.

7. What is a common symptom of stroke?

- A. Severe headache
- B. Sudden numbness or weakness of the face, arm, or leg**
- C. Chest pain
- D. Difficulty breathing

Sudden numbness or weakness of the face, arm, or leg is a hallmark symptom of a stroke. This sign is often characterized by the rapid onset of weakness or loss of control in one side of the body, typically affecting the arm, leg, or facial muscles. This is due to the interruption of blood flow to the brain, which can be caused by either an ischemic stroke (where blood flow is blocked) or a hemorrhagic stroke (where blood vessels burst). Recognizing this symptom is crucial for timely medical intervention, as the likelihood of better outcomes increases significantly the sooner treatment begins. Severe headache is indeed associated with certain types of strokes, particularly hemorrhagic strokes but is less common in ischemic strokes. Chest pain and difficulty breathing are symptoms associated with other medical conditions, such as cardiac events, and are not typical presentations of stroke. While they can occur in a stroke patient or due to other concurrent health issues, they do not represent core stroke diagnostic criteria. Understanding the distinct symptoms of a stroke can facilitate immediate recognition and treatment, which is critical in improving patient prognosis.

8. Which change in speech may indicate a patient has had a stroke?

- A. Clear speech with normal volume
- B. Slurred speech or difficulty finding words**
- C. Excessive talking
- D. Sudden increase in vocabulary

The identification of slurred speech or difficulty finding words as an indicator of a potential stroke is rooted in the understanding of how strokes affect brain function, particularly in areas associated with language. When a stroke occurs, it can disrupt blood flow to specific regions in the brain responsible for speech production and comprehension, such as Broca's area and Wernicke's area. Patients experiencing a stroke may present with dysarthria, where the muscles used for speech are weakened, leading to slurred or unclear speech. Additionally, they might struggle to articulate thoughts or find the correct words to express themselves, known as aphasia. These speech changes are critical symptoms of stroke and warrant immediate medical attention, as they can help healthcare professionals assess the severity and location of neurological impairment. Other options like clear speech with normal volume do not indicate a stroke, as this is a sign of intact speech capabilities. Excessive talking could be related to various behavioral or psychological states, rather than an indication of neurological damage. A sudden increase in vocabulary is also not consistent with stroke symptoms and may suggest improvement or adaptation in communication rather than a sign of impairment. Therefore, slurred speech or difficulty finding words serves as a significant indicator for the recognition of a stroke and the need for urgent intervention

9. Which condition is primarily indicated by Wallenberg syndrome?

- A. Basilar artery occlusion
- B. PICA stroke**
- C. MCA stroke
- D. ACA stroke

Wallenberg syndrome, also known as lateral medullary syndrome, is primarily associated with a stroke in the region supplied by the posterior inferior cerebellar artery (PICA). This syndrome results from an impairment of blood flow to the brainstem, particularly affecting the medulla oblongata. The core features of Wallenberg syndrome include symptoms such as dysphagia, hoarseness, and ataxia, which arise from lesions in specific cranial nerve and brainstem pathways located in the lateral medulla. Blood supply to this area is predominantly from PICA, making it the most likely artery involved in Wallenberg syndrome. In contrast, the other options refer to strokes affecting different vascular territories that do not typically result in the signs and symptoms characteristic of Wallenberg syndrome. For instance, strokes in the basilar artery, middle cerebral artery (MCA), or anterior cerebral artery (ACA) would manifest with distinct clinical presentations attributed to their respective regions, not the unique combination seen in Wallenberg syndrome.

10. How can caregivers best support stroke recovery?

- A. By minimizing patient interactions
- B. By providing encouragement and assistance with therapy exercises**
- C. By discouraging any physical activity
- D. By focusing solely on medication management

Providing encouragement and assistance with therapy exercises is vital for stroke recovery because it significantly impacts a patient's overall rehabilitation process. Stroke can lead to physical and cognitive impairments, making it essential for caregivers to engage patients actively in their recovery. Active participation in therapy exercises helps rebuild strength, coordination, and functional independence, which are critical for regaining skills lost due to the stroke. Furthermore, encouragement fosters a positive environment and can boost the patient's motivation and self-esteem. It promotes adherence to therapy regimens, as patients are more likely to engage in exercises when they feel supported and understood. This supportive approach allows caregivers to build a collaborative relationship that enhances recovery outcomes and provides emotional support necessary for coping with the challenges of rehabilitation.

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

<https://strokecertifiedregisterednurse.examzify.com>

We wish you the very best on your exam journey. You've got this!

SAMPLE